

## IIa. Systems of many electrons

---

### Exchange of particle labels

Consider a system of  $n$  identical particles. By identical particles we mean that all intrinsic properties of the particles are the same, such as mass, spin, charge, etc.

*Figure II.1 A pairwise permutation of the labels on identical particles.*

In classical mechanics we can in principle follow the trajectory of each individual particle. They are, therefore, distinguishable even though they are identical. In quantum mechanics the particles are not distinguishable if they are close enough or if they interact strongly enough. The quantum mechanical wave function must reflect this fact.

When the particles are indistinguishable, the act of labeling the particles is an arbitrary operation without physical significance. Therefore, all observables must be unaffected by interchange of particle labels. The operators are said to be symmetric under interchange of labels. The Hamiltonian, for example, is symmetric, since the intrinsic properties of all the particles are the same. Let  $\psi(1, 2, \dots, n)$  be a solution to the Schrödinger equation, (Here  $n$  represents all the coordinates, both spatial and spin, of the particle labeled with  $n$ ). Let  $P_{ij}$  be an operator that permutes (or ‘exchanges’) the labels  $i$  and  $j$ :

$$P_{ij}\psi(1, 2, \dots, i, \dots, j, \dots, n) = \psi(1, 2, \dots, j, \dots, i, \dots, n).$$

This means that the function  $P_{ij}\psi$  depends on the coordinates of particle  $j$  in the same way that  $\psi$  depends on the coordinates of particle  $i$ . Since  $H$  is symmetric under interchange of labels:

$$H(P_{ij}\psi) = P_{ij}H\psi, \quad \text{i.e.,} \quad [P_{ij}, H] = 0.$$

Therefore,  $P_{ij}\psi$  is also a solution of the Schrödinger equation with the same eigenvalue as  $\psi$ .

*Exchange Degeneracy:* There are  $n!$  different permutations of  $n$  labels and all the  $n!$  wave functions have the same energy. Some linear combination of the  $n!$  functions gives the proper description of the system. Since  $[P_{ij}, H] = 0$ , i.e.  $P_{ij}$  is a constant of the motion,

the linear combination that describes the system initially, is the proper linear combination for all time.

The density (i.e. probability distribution) must be unaffected by  $P$

$$|P_{ij}\psi|^2 = |\psi|^2$$

but that is not the case with the wavefunction, which is not an observable. Therefore, the functions  $\psi$  and  $P_{ij}\psi$  can differ in phase

$$P_{ij}\psi = e^{i\alpha}\psi$$

where  $\alpha$  is a real number. Applying  $P_{ij}$  again will undo the permutation, so we must have

$$P_{ij}(P_{ij}\psi) = P_{ij}^2\psi = \psi$$

that is

$$(e^{i\alpha})^2 = 1.$$

There are only two distinct solutions  $\alpha = 0$  and  $\alpha = \pi$  corresponding to

$$P_{ij}\psi = \begin{cases} +\psi & \text{symmetric under interchange} \\ -\psi & \text{antisymmetric under interchange} \end{cases} .$$

It turns out that both solutions are found in nature. For some particles, called *Fermions*, the wavefunction should be *antisymmetric* ( $\alpha = \pi$ ). Examples are electrons, protons and neutrons. For other particles, called *Bosons*, the wavefunction should be *symmetric* ( $\alpha = 0$ ). Examples are photons (particles with integral spin).

The behavior of the wavefunction when composite particles (not elementary), such as atoms, are interchanged can be deduced by counting the number of Fermions. For example: When two  ${}^4\text{He}$  atoms are interchanged

2 Electrons:	Pick up factor	$(-1)(-1) = +1$
2 Protons:	Pick up factor	$(-1)(-1) = +1$
2 Neutrons:	Pick up factor	$(-1)(-1) = +1$ .

So the wavefunction should not change sign, i.e.  ${}^4\text{He}$  atoms behave as Bosons.

When two  ${}^3\text{He}$  atoms are interchanged:

2 Electrons:	Pick up factor	$(-1)(-1) = +1$
2 Protons:	Pick up factor	$(-1)(-1) = +1$
1 Neutron:	Pick up factor	$(-1)$ .

So the wavefunction should change sign,  ${}^3\text{He}$  atoms are Fermions.

Since hydrogen atoms are very light, one often needs to use a quantum mechanical description of the motion of the nuclei as well as the electrons. The wave function changes sign if we interchange labels of two electrons (electrons are Fermions) and it changes sign if we interchange the labels of two nuclei (the nuclei consist of one proton which is a

Fermion), but if we interchange two  $H$  atoms (both electron and proton), then we pick up a factor of  $(-1)(-1) = +1$ , i.e., the wavefunction does not change sign. Hydrogen atoms behave as Bosons.

When a stationary state wave function is being constructed for a system, for example to evaluate possible values of the energy, it is not enough to just solve for eigenfunctions of the Hamiltonian operator, one also needs to build in the right symmetry with respect to the exchange of labels on the particles. For example, if a given a function  $\psi(1, 2, \dots, n)$  is found to satisfy the Schrödinger equation but does not have the right exchange symmetry, it is necessary to construct a linear combination of permutations of the labels to obtain a function that is symmetric under the exchange of labels when dealing with Bosons or antisymmetric when dealing with Fermions. Let  $P$  be any one of the  $n!$  permutations. It can be constructed from a sequence of pairwise permutations. Let the number of pairwise permutations required be  $r_P$ . For example:

$$\begin{array}{rcll}
 & (1, 3, 2) & r = 1 & \text{odd} \\
 & \downarrow P_{13} & & \\
 (1, 2, 3) & (2, 3, 1) & r = 2 & \text{even} \\
 & \downarrow P_{12} & & \\
 & (2, 1, 3) & \left\{ \begin{array}{l} r = 3 \\ \text{or } r = 1 \end{array} \right. & \text{odd}
 \end{array}$$

The normalized symmetric linear combination appropriate for Bosons is

$$\psi^+ = \frac{1}{\sqrt{n!}} \sum_p P_p \phi(1, 2, \dots, n) .$$

The index  $p$  runs over all the  $n!$  permutations. Similarly, the normalized antisymmetric linear combination appropriate for Fermions is:

$$\psi^- = \frac{1}{\sqrt{n!}} \sum_p (-1)^{r_p} P_p \phi(1, 2, \dots, n) .$$

The difference in the energy that is obtained from  $\phi$  and that obtained from  $\psi^-$  is called the *exchange energy*. That is

$$E_{exch} = \langle \psi^- | H | \psi^- \rangle - \langle \phi | H | \phi \rangle .$$

We can define operators that perform the required linear combination. The *symmetrizer*,  $S$ , is

$$S \equiv \frac{1}{\sqrt{n!}} \sum P$$

and the *antisymmetrizer*,  $A$ , is

$$A \equiv \frac{1}{\sqrt{n!}} \sum (-1)^{r_p} P .$$

Then  $\psi^+ = S\phi$  and  $\psi^- = A\phi$ .

**Non-Interacting Electrons:** (the simplest system containing several electrons)

The total Hamiltonian of the system is a sum of terms, each acting only on a single electron:

$$H = H_1 + H_2 + \dots H_n .$$

Since the Hamiltonian separates, a solution to the Schrödinger equation can be found that is a product of one electron functions:

$$\phi(1, 2, \dots, n) = \chi_a(1)\chi_b(2) \dots \chi_0(n) .$$

The one electron functions are called *spin-orbitals* and depend both on spin and spatial coordinates and the product function,  $\chi(1, 2, \dots, n)$ , is called a *Hartree Product*. The properly antisymmetrized many electron wavefunction is:

$$\begin{aligned} \psi^-(1, 2, \dots, n) &= A \phi(1, 2, \dots, n) \\ &= \frac{1}{\sqrt{n!}} \begin{vmatrix} \chi_a(1) & \chi_b(1) & \dots & \chi_0(1) \\ \chi_a(2) & \chi_b(2) & \dots & \chi_0(2) \\ \vdots & \vdots & \vdots & \vdots \\ \chi_a(n) & \chi_b(n) & \dots & \chi_0(n) \end{vmatrix} . \end{aligned}$$

Applying some general rules about determinants, we can see that this wavefunction has the required properties: When the determinant is expanded out there are  $n!$  terms half of which have a minus sign. A permutation  $P_{ij}$  corresponds to interchanging the rows  $i$  and  $j$ , which causes the determinant to change sign. Therefore the antisymmetry is built in. We will frequently be dealing with determinantal wave functions in this section. It is therefore essential to have a convenient short hand notation for such functions. We will use the notation

$$\psi^-(1, 2, \dots, n) = |\chi_a(1)\chi_b(2) \dots \chi_0(n) \rangle .$$

The pointed bracket will therefore have a different meaning in this section than it had previously.

If two spin orbitals are equal, for example  $\chi_a = \chi_b$  then two columns are the same and the determinant vanishes,  $\psi = 0$ . That is, two electrons cannot be in the same spin-orbital. This is referred to as *Pauli exclusion*.

Assuming the hamiltonian does not couple spatial coordinates and spin, the spin-orbitals can be written as a product of spatial and spin functions. For example, if electron 1 is in spatial orbital  $a$  and has spin up, the spin-orbital is

$$\chi_a(1) = \phi_I(\vec{r}_1)\alpha(\omega_1) .$$

Since we have used the symbols of the ket notation to mean a determinantal wave function in this section, we will be explicitly dealing with functions rather than kets. To deal with the spin, we therefore imagine having functions  $\alpha(\omega)$  for spin up and  $\beta(\omega)$  for spin down and introduce, purely for convenience, a hypothetical variable,  $\omega$ . Eventually, we will always integrate over expressions involving the functions  $\alpha$  and  $\beta$  and the important thing to remember is that they are orthonormal

$$\int d\omega \alpha^*(\omega) \beta(\omega) = 0$$

and

$$\int d\omega \alpha^*(\omega) \alpha(\omega) = 1.$$

The Hartree product is truly an independent electron wavefunction. But the antisymmetrized wavefunction obtained by constructing the Slater determinant has introduced some, limited correlation between any pair of electrons that have parallel spin. This is called the *exchange correlation* (not to be confused with the 'correlation energy' which is the difference between the Hartree-Fock estimate of the energy and the exact value of the energy). The effect can be seen by looking at the electron density in configuration space. Consider a two electron Slater determinant:

$$\psi^-(1, 2) = A \chi_a(1)\chi_b(2) = |\chi_a(1)\chi_b(2) \rangle .$$

*Example a.* First, assume the two electrons have opposite spins and occupy different spatial orbitals

$$\chi_a(1) = \phi_I(\vec{r}_1)\alpha(\omega_1)$$

and

$$\chi_b(2) = \phi_{II}(\vec{r}_2) \beta(\omega_2) .$$

( $\alpha$  means spin up and  $\beta$  spin down). By expanding the determinant, we get:

$$\psi^-(1, 2) = \frac{1}{2} (\phi_I(\vec{r}_1)\alpha(\omega_1)\phi_{II}(\vec{r}_2)\beta(\omega_2) - \phi_I(\vec{r}_2)\alpha(\omega_2)\phi_{II}(\vec{r}_1)\beta(\omega_1)) .$$

The simultaneous probability of finding one of the two electron near a point  $\vec{r}_s$  and the other near a point  $\vec{r}_t$  (that is, one of the electrons is in an interval between  $\vec{r}_s$  and  $\vec{r}_s + d\vec{r}$  and the other electron in an interval between  $\vec{r}_t$  and  $\vec{r}_t + d\vec{r}$ ) is obtained from the magnitude squared of the wave function  $|\psi(1, 2)|^2$  by integrating out the 'spin variables':

$$\begin{aligned} P(\vec{r}_1, \vec{r}_2)d\vec{r}_1d\vec{r}_2 &= \int d\omega_1 \int d\omega_2 |\psi(1, 2)|^2 d\vec{r}_1 d\vec{r}_2 \\ &= \frac{1}{2} \left[ |\phi_I(\vec{r}_1)|^2 |\phi_{II}(\vec{r}_2)|^2 + |\phi_I(\vec{r}_2)|^2 |\phi_{II}(\vec{r}_1)|^2 \right] d\vec{r}_1 d\vec{r}_2 . \end{aligned}$$

By setting  $\vec{r}_1 = \vec{r}_s$  and  $\vec{r}_2 = \vec{r}_t$ , the first term is the product of the probability of finding electron 1 within  $\vec{r}_s$  and  $(\vec{r}_s + d\vec{r})$  times the probability of finding electron 2 within  $\vec{r}_t$  and

$(\vec{r}_t + d\vec{r})$  as electron 1 occupies orbital  $\phi_I$  and electron 2 occupies orbital  $\phi_{II}$ . The second term has electron 2 occupying  $\phi_I$  and electron 1 occupying  $\phi_{II}$ . Since the electrons are indistinguishable, the correct probability is the average of the two terms. Therefore, two electrons with opposite spin are fully uncorrelated at this level of theory.

*Example b.* Secondly, assume the two electrons have the same spin, say spin down. Then

$$\begin{aligned}\chi_a(1) &= \phi_I(\vec{r}_1)\beta(\omega_1) \\ \chi_b(2) &= \phi_{II}(\vec{r}_2)\beta(\omega_2) .\end{aligned}$$

Then, the probability distribution becomes:

$$\begin{aligned}P(\vec{r}_1, \vec{r}_2) &= \frac{1}{2} \left[ |\phi_I(\vec{r}_1)|^2 |\phi_{II}(\vec{r}_2)|^2 + |\phi_I(\vec{r}_2)|^2 |\phi_{II}(\vec{r}_1)|^2 \right. \\ &\quad \left. - 2\text{Re} \left\{ \phi_I^*(\vec{r}_1) \phi_{II}(\vec{r}_1) \phi_{II}^*(\vec{r}_2) \phi_I(\vec{r}_2) \right\} \right] .\end{aligned}$$

The extra term that has appeared as compared with example *a* introduces *correlation* in the distribution of the two electrons. In particular  $P(\vec{r}_1 = \vec{r}, \vec{r}_2 = \vec{r}) = 0$ , i.e., two electrons with parallel spin cannot be found at the same point in space. A *Fermi hole* is said to exist around each electron, a region from which other electrons of the same spin are excluded. The presence of the Fermi hole is an important consideration. It results, for example, in a substantial difference in the accuracy of the Hartree-Fock approximation for singlet states and for triplet states, since the presence of the extra Fermi hole in the triplet tends to keep the electrons apart and, thereby, lower the Coulomb repulsion between them.

### Calculation of Matrix Elements using Slater Determinants:

Let  $\psi(1, 2, 3, \dots, N)$  be a normalized Slater determinant

$$\psi(1, 2, \dots, N) = \frac{1}{\sqrt{N!}} \begin{vmatrix} \chi_i(1) & \chi_j(1) & \dots & \chi_k(1) \\ \chi_i(2) & \chi_j(2) & \dots & \chi_k(2) \\ \vdots & \vdots & \ddots & \vdots \\ \chi_i(N) & \chi_j(N) & \dots & \chi_k(N) \end{vmatrix}$$

where the  $\chi_i$  are spin-orbitals. We will use the shorthand notation

$$\psi(1, 2, \dots, N) = |\chi_i \chi_j \dots \chi_k \rangle$$

where it is understood that the electron labels are in the order  $1, 2, \dots, N$ . Because of the antisymmetry we have, for example:

$$|\dots \chi_m \dots \chi_n \dots \rangle = -|\dots \chi_n \dots \chi_m \dots \rangle .$$

Various calculations in quantum mechanics involve the evaluation of matrix elements of an operator with the wavefunction. It, therefore, becomes important to develop expressions for matrix elements of various types of operators when the wavefunction is of the form of

a Slater determinant. Given an operator  $O$  and two  $N$ -electron determinants  $|K\rangle$  and  $|L\rangle$ , the problem now is to evaluate the matrix element  $\langle K|O|L\rangle$ . We need to obtain an expression involving integrals over the individual spin-orbitals. In particular, we need to evaluate the energy,  $\langle H\rangle = \langle K|H|L\rangle$ .

There are two types of operators that are relevant:

*One electron operators* are of the form

$$O_1 = \sum_{i=1}^N h(i)$$

where  $h(i)$  only involves the  $i$ -th electron and the summation index,  $i$ , runs over all electrons. For example

$$h(i) = -\frac{\hbar^2}{2m} \nabla_i^2.$$

*Two electron operators* are of the form

$$O_2 = \sum_{i=1}^N \sum_{j>i}^N v(i, j) \equiv \sum_{i<j} v(i, j)$$

where  $v(i, j)$  is an operator that depends on the coordinates of both electrons,  $i$  and  $j$ . An important example is the Coulomb interaction  $e^2/r_{ij}$  and we will be using that example in the following discussion.

Regarding the determinants  $|K\rangle$  and  $|L\rangle$ , there are three different situations:

*Case 1:* The two determinants are equal, i.e.,

$$|L\rangle = |K\rangle = |\dots \chi_m \chi_n \dots\rangle.$$

*Case 2:* The two determinants differ by one spin-orbital: For example, replacing  $\chi_m$  by  $\chi_p$  in  $|L\rangle$

$$\begin{aligned} |K\rangle &= |\dots \chi_m \chi_n \dots\rangle \\ |L\rangle &= |\dots \chi_p \chi_n \dots\rangle. \end{aligned}$$

*Case 3:* The two determinants differ by two spin-orbitals: For example, replacing  $\chi_m$  by  $\chi_p$  and  $\chi_n$  by  $\chi_q$  in  $|L\rangle$

$$\begin{aligned} |K\rangle &= |\dots \chi_m \chi_n \dots\rangle \\ |L\rangle &= |\dots \chi_p \chi_q \dots\rangle. \end{aligned}$$

It can easily be shown that the matrix elements of  $O_1$  are zero if  $|K\rangle$  and  $|L\rangle$  differ by more than one spin-orbital, and that the matrix elements of  $O_2$  are zero if  $|K\rangle$  and  $|L\rangle$  differ by more than two spin-orbitals.

It is essential to use some convenient shorthand notation for the various integrals. Unfortunately, there are two different notations in common use, that are sometimes referred

to as ‘chemists notation’ and ‘physicists notation’. We will use the former and denote integrals over spin-orbitals with square brackets, [ ]. The definitions are as follows:

$$[i|h|j] \equiv \int dx_1 \chi_i^*(1) h(\vec{r}_1) \chi_j(1)$$

and

$$[ij|k\ell] \equiv \int dx_1 \int dx_2 \chi_i^*(1) \chi_j(1) \frac{1}{r_{12}} \chi_k^*(2) \chi_\ell(2)$$

The integration variable  $x_i$  denotes both the spatial and spin coordinates of electron  $i$  (note, we assume the Hamiltonian does not depend on spin and therefore denote the variable there as  $\vec{r}_1$ ).

The following table gives the matrix elements of *one-electron* operators for determinant wave functions in terms of the integrals over spin-orbitals.

Table 1: Matrix elements of one electron operators of the form  $O_1 = \sum_{i=1}^N h(i)$

---

Case 1:  $|K\rangle = |\dots mn\dots\rangle$

$$\langle K|O_1|K\rangle = \sum_m^N [m|h|m]$$

Case 2:  $|K\rangle = |\dots mn\dots\rangle$

$|L\rangle = |\dots pn\dots\rangle$

$$\langle K|O_1|L\rangle = [m|h|p]$$

Case 3:  $|K\rangle = |\dots mn\dots\rangle$

$|L\rangle = |\dots pq\dots\rangle$

$$\langle K|O_1|L\rangle = 0$$

---



The following table gives the matrix elements of the *two-electron* operator representing Coulomb interaction of  $N$  electrons in terms of the spin-orbitals of the determinants.

Table 2: Matrix elements of two electron operators of the form  $O_2 = \sum_{i=1}^N \sum_{j>i}^N r_{ij}^{-1}$

---

Case 1:  $|K\rangle = |\dots nm\dots\rangle$

$$\langle K|O_2|K\rangle = \frac{1}{2} \sum_m^N \sum_n^N [mm|nn] - [mn|nm]$$

Case 2:  $|K\rangle = |\dots mn\dots\rangle$

$|L\rangle = |\dots pn\dots\rangle$

$$\langle K|O_2|L\rangle = \sum_n^N [mp|nn] - [mn|np]$$

Case 3:  $|K\rangle = |\dots mn\dots\rangle$

$|L\rangle = |\dots pq\dots\rangle$

$$\langle K|O_2|L\rangle = [mp|nq] - [mq|np]$$


---

To illustrate how the rules in the table come about, calculations of some of the matrix elements are presented below for some of the possible choices of Slater determinants.

*Example 1:* The simplest case is when the same Slater determinant appears on the left and right hand side of the matrix element (case 1)

$$|\psi\rangle = |\chi_1\chi_2\rangle$$

1.a For a one-electron operator:

$$\begin{aligned} \langle \psi|h(1)|\psi\rangle &= \int d\vec{x}_1 \int d\vec{x}_2 \frac{1}{\sqrt{2}} \left( \chi_1(\vec{x}_1)\chi_2(\vec{x}_2) - \chi_2(\vec{x}_1)\chi_1(\vec{x}_2) \right)^* \\ &\quad h(\vec{r}_1) \frac{1}{\sqrt{2}} \left( \chi_1(\vec{x}_1)\chi_2(\vec{x}_2) - \chi_2(\vec{x}_1)\chi_1(\vec{x}_2) \right) \\ &= \frac{1}{2} \int d\vec{x}_1 \int d\vec{x}_2 \left( \chi_1^*(\vec{x}_1)\chi_2^*(\vec{x}_2)h(\vec{r}_1)\chi_1(\vec{x}_1)\chi_2(\vec{x}_2) \right. \\ &\quad + \chi_2^*(\vec{x}_1)\chi_1^*(\vec{x}_2)h(\vec{r}_1)\chi_2(\vec{x}_1)\chi_1(\vec{x}_2) \\ &\quad - \chi_1^*(\vec{x}_1)\chi_2^*(\vec{x}_2)h(\vec{r}_1)\chi_2(\vec{x}_1)\chi_1(\vec{x}_2) \\ &\quad \left. - \chi_2^*(\vec{x}_1)\chi_1^*(\vec{x}_2)h(\vec{r}_1)\chi_1(\vec{x}_1)\chi_2(\vec{x}_2) \right). \end{aligned}$$

Since the operator  $h(1)$  does not involve coordinates of electron 2 we can easily carry out the  $\vec{x}_2$  integration. In the first two terms,  $\vec{x}_2$  only appears in the same spin-orbital within each term. Therefore using the fact that both  $\chi_1$  and  $\chi_2$  are normalized, the integration over  $\vec{x}_2$  gives 1 in the first two terms. However, in the last two terms,  $\vec{x}_2$  appears in both  $\chi_1$  and  $\chi_2$  within each term. The integration over  $\vec{x}_2$  then gives zero because the spin-orbitals are orthogonal. We are left with

$$\begin{aligned} \langle \psi | h(1) | \psi \rangle &= \frac{1}{2} \int d\vec{x}_1 \chi_1^*(\vec{x}_1) h(\vec{r}_1) \chi_1(\vec{x}_1) \\ &\quad + \frac{1}{2} \int d\vec{x}_1 \chi_2^*(\vec{x}_1) h(\vec{r}_1) \chi_1^*(\vec{x}_2) \\ &= \frac{1}{2} \sum_m^2 [m|h|m]. \end{aligned}$$

Similarly

$$\langle \psi | h(2) | \psi \rangle = \frac{1}{2} \sum_m^2 [m|h|m].$$

Adding the two gives case 1 in the table for the one-electron operator,  $O_1$ .

1.b For the two-electron operator  $O_2 = 1/r_{12}$ :

$$\begin{aligned} \langle \psi | \frac{1}{r_{12}} | \psi \rangle &= \frac{1}{2} \int d\vec{x}_1 \int d\vec{x}_2 \left( \chi_1^*(\vec{x}_1) \chi_2^*(\vec{x}_2) \frac{1}{r_{12}} \chi_1(\vec{x}_1) \chi_2(\vec{x}_2) \right. \\ &\quad + \chi_2^*(\vec{x}_1) \chi_1^*(\vec{x}_2) \frac{1}{r_{12}} \chi_2(\vec{x}_1) \chi_1(\vec{x}_2) \\ &\quad - \chi_1^*(\vec{x}_1) \chi_2^*(\vec{x}_2) \frac{1}{r_{12}} \chi_2(\vec{x}_1) \chi_1(\vec{x}_2) \\ &\quad \left. - \chi_2^*(\vec{x}_1) \chi_1^*(\vec{x}_2) \frac{1}{r_{12}} \chi_1(\vec{x}_1) \chi_2(\vec{x}_2) \right) \end{aligned}$$

Since  $1/r_{12} = 1/r_{21}$ , the indices can be interchanged on the dummy integration variables. Therefore, the first term is the same as the second and the third term is the same as the fourth.

$$\langle \psi | \frac{1}{r_{12}} | \psi \rangle = [11|22] - [12|21].$$

This illustrates case 1 in the table for the two-electron operator,  $O_2$ .

*Example 2:* Now use determinants that differ by one spin-orbital (case 3). Again, the Slater determinants are constructed by using two spin-orbitals but different spin-orbitals are chosen for the two determinants:

$$|\psi_A \rangle = |\chi_1 \chi_2 \rangle$$

and

$$|\psi_B \rangle = |\chi_3 \chi_4 \rangle .$$

We clearly have

$$\langle \psi_A | h(1) | \psi_B \rangle = 0$$

because the  $\vec{x}_2$  integration gives zero in all terms,

$$0 = \int d\vec{x}_2 \chi_i(\vec{x}_2) \chi_j(\vec{x}_2) \text{ when } i \neq j.$$

This illustrates case 3 in the table for the one electron operator,  $O_1$ .

The two-electron matrix element is, by definition,

$$\langle \psi_A | \frac{1}{r_{12}} | \psi_B \rangle = [13|24] - [14|23].$$

*The energy of a determinantal wavefunction:*

Using the tables, we can readily find the expectation value of the total energy in a single determinant state  $|K \rangle$  of  $N$  electrons

$$\begin{aligned} \langle H \rangle &= \langle K | H | K \rangle = \langle K | O_1 + O_2 | K \rangle \\ &= \sum_m^N [m|h|m] + \frac{1}{2} \sum_m^N \sum_n^N [mm|nn] - [mn|nm] \end{aligned}$$

The sums run over all the occupied spin-orbitals. The one electron operator  $h$  includes the kinetic energy of the electron and the interaction with all the nuclei

$$h(i) = -\frac{1}{2} \nabla_i^2 - \sum_A \frac{Z_A}{r_{iA}}$$

The summation index  $A$  runs over all nuclei that electron  $i$  interacts with. Here the electron charge has been set to one,  $e = 1$ , and  $Z_A$  is the nuclear charge in units of the electron charge,  $e$ . Each occupied spin-orbital  $\chi_i$  in  $|K \rangle$  contributes  $[i|h|i]$  to the energy and every unique pair of occupied spin-orbitals  $\chi_i, \chi_j$  contributes  $\langle ij||ij \rangle$ . Note that

$$\frac{1}{2} \sum_m^N \sum_n^N [mm|nn] - [mn|nm] = \sum_m \sum_{n>m} [mm|nn] - [mn|nm]$$

since

$$[mn|nm] = [nm|mn].$$

and terms with  $n = m$  cancel out.